

BEST AVAILABLE COPY

IN THE CLAIMS

Claims 1-4 (canceled)

Claim 5 (currently amended): A method of enhancing wound healing in an external wound of an individual, comprising the ~~step~~ steps of:

covering said wound with a wound coverage material, wherein said wound coverage material is impregnated with a therapeutically effective amount of a cholesterol-containing cationic liposome, said liposome comprising at least one gene construct encoding a growth factor.

Claim 6 (original): The method of claim 5, wherein said impregnation of said wound coverage material is performed prior to covering said wound or subsequent to covering said wound.

Claim 7 (original): The method of claim 5, wherein said wound is selected from the group consisting of thermal trauma, chemical trauma, excisional trauma, surgical trauma and abrasion.

Claim 8 (original): The method of claim 5, wherein said wound coverage material is selected from the group consisting of human fetal amnion, human fetal chorion, human cadaver skin and synthetic skin.

Claim 9 (original): The method of claim 5, wherein said growth factor is selected from the group consisting of growth hormone, insulin-like growth factor-I, keratinocyte growth factor, fibroblast growth factor, epidermal growth factor, platelet derived growth factor and transforming growth factor- β .

Claim 10 (original): The method of claim 9, wherein said growth factor is insulin-like growth factor-I (IGF-I) and the concentration of said gene encoding IGF-I in the liposomes is about 2.2 $\mu\text{g}/10 \mu\text{l}$ liposomes.

Claim 11 (currently amended): A method of enhancing wound healing in an external wound of an individual, comprising the ~~step~~ steps of:

BEST AVAILABLE COPY

covering said wound with a wound closure material, wherein said wound closure material is impregnated with a therapeutically effective amount of a cholesterol-containing cationic liposome, said liposome comprising at least one gene construct encoding a growth factor.

Claim 12 (currently amended): The method of claim 11, wherein said impregnation of said wound ~~coverage~~ closure material is performed prior to covering said wound or subsequent to covering said wound.

Claim 13 (original): The method of claim 11, wherein said wound is selected from the group consisting of thermal trauma, chemical trauma, excisional trauma, surgical trauma and abrasion.

Claim 14 (original): The method of claim 11, wherein said wound closure material is selected from the group consisting of human fetal amnion, human fetal chorion, human syngeneic skin, and human allogeneic skin.

Best Available Copy

Claim 15 (original): The method of claim 11, wherein said growth factor is selected from the group consisting of growth hormone, insulin-like growth factor-I, keratinocyte growth factor, fibroblast growth factor, epidermal growth factor, platelet derived growth factor and transforming growth factor- β .

Claim 16 (original): The method of claim 15, wherein said growth factor is insulin-like growth factor-I (IGF-I) and the concentration of said gene encoding IGF-I in the liposomes is about 2.2 $\mu\text{g}/10 \mu\text{l}$ liposomes.